

## INTERVENTIONAL IMAGING PREP

<p><b>Abdominal Aorta w/Runoff</b></p>	<p>Please call 982-3477 to schedule a pre-admission appointment.  <b>Nothing to eat or drink for 6 hours prior to exam</b> (can have daily medications with a sip of water). <b>Withhold Coumadin for 5 days, Lovenox for 1 dose, Plavix for 5 days, Pletal for 4 days, Pradaxa for 48 hours, &amp; Ticlid for 14 days prior to exam after obtaining approval from prescribing physician.</b> Lab work as indicated at time of scheduling. Must have a <b>driver</b>.</p>
<p><b>Abdominal Aortogram</b></p>	<p>Please call 982-3477 to schedule a pre-admission appointment.  <b>Nothing to eat or drink for 6 hours prior to exam</b> (can have daily medications with a sip of water). <b>Withhold Coumadin for 5 days, Lovenox for 1 dose, Plavix for 5 days, Pletal for 4 days, Pradaxa for 48 hours, &amp; Ticlid for 14 days prior to exam after obtaining approval from prescribing physician.</b> Lab work as indicated at time of scheduling. Must have a <b>driver</b>.</p>
<p><b>Abscess Drainage</b></p>	<p>Please call 982-3477 to schedule a pre-admission appointment.  <b>Nothing to eat or drink for 6 hours prior to exam</b> (can have daily medications with a sip of water). <b>Withhold Coumadin for 5 days &amp; Lovenox for 1 dose prior to exam after obtaining approval from prescribing physician.</b> Lab work as indicated at time of scheduling. Must have a <b>driver</b>.</p>
<p><b>Abscess-O-Gram</b></p>	<p>No Prep, unless instructed otherwise</p>
<p><b>Arthrogram, Wrist</b></p>	<p>No Prep, unless instructed otherwise</p>
<p><b>Biliary Catheter Check</b></p>	<p>No Prep, unless instructed otherwise</p>
<p><b>Biliary Catheter Exchange</b></p>	<p>Please call 982-3477 to schedule a pre-admission appointment.  <b>Nothing to eat or drink for 6 hours prior to exam</b> (can have daily medications with a sip of water). <b>Withhold Coumadin for 5 days &amp; Lovenox for 1 dose prior to exam after obtaining approval from prescribing physician.</b> Lab work as indicated at time of scheduling. Must have a <b>driver</b>.</p>
<p><b>Bone Biopsy-Deep</b></p>	<p>Please call 982-3477 to schedule a pre-admission appointment.  <b>Nothing to eat or drink for 6 hours prior to exam</b> (can have daily medications with a sip of water). <b>Nothing to eat or drink for 6 hours prior to exam</b> (can have daily medications with a sip of water). <b>Withhold Coumadin for 5 days, Lovenox for 1 dose, Plavix for 5 days, Pletal for 4 days, Pradaxa for 48 hours, &amp; Ticlid for 14 days prior to exam after obtaining approval from prescribing physician.</b> Lab work as indicated at time of scheduling. Must have a <b>driver</b>.</p>
<p><b>Carotid Cerebral</b></p>	<p>Please call 982-3477 to schedule a pre-admission</p>

<p><b>Angiogram/Arteriogram</b></p>	<p>appointment.  <b>Nothing to eat or drink for 6 hours prior to exam</b> (can have daily medications with a sip of water. <b>Withhold Coumadin for 5 days, Lovenox for 1 dose, Plavix for 5 days, Pletal for 4 days, Pradaxa for 48 hours, &amp; Ticlid for 14 days prior to exam after obtaining approval from prescribing physician.</b> Lab work as indicated at time of scheduling. (<b>Anesthesia patients</b> must have nothing to eat or drink for 8 hours prior to exam.) Must have a <b>driver</b>.</p>
<p><b>Carotid Cervical Angiogram/Arteriogram</b></p>	<p>Please call 982-3477 to schedule a pre-admission appointment.  <b>Nothing to eat or drink for 6 hours prior to exam</b> (can have daily medications with a sip of water. <b>Withhold Coumadin for 5 days, Lovenox for 1 dose, Plavix for 5 days, Pletal for 4 days, Pradaxa for 48 hours, &amp; Ticlid for 14 days prior to exam after obtaining approval from prescribing physician.</b> Lab work as indicated at time of scheduling. (<b>Anesthesia patients</b> must have nothing to eat or drink for 8 hours prior to exam.) Must have a <b>driver</b>.</p>
<p><b>Chest Tube Placement</b></p>	<p>Please call 982-3477 to schedule a pre-admission appointment.  <b>Nothing to eat or drink for 6 hours prior to exam</b> (can have daily medications with a sip of water. <b>Withhold Coumadin for 5 days, Lovenox for 1 dose, Plavix for 5 days, Pletal for 4 days, Pradaxa for 48 hours, &amp; Ticlid for 14 days prior to exam after obtaining approval from prescribing physician.</b> Lab work as indicated at time of scheduling. Must have a <b>driver</b>.</p>
<p><b>Cisternogram</b></p>	<p>Please call 982-3477 to schedule a pre-admission appointment.  <b>Nothing to eat or drink for 6 hours prior</b> (can have daily medications with a sip of water. <b>Withhold Coumadin for 5 days, Lovenox for 1 dose, Plavix for 5 days, Pletal for 4 days, Pradaxa for 48 hours, &amp; Ticlid for 14 days prior to exam after obtaining approval from prescribing physician.</b> Lab work as indicated at time of scheduling. Must have a <b>driver</b>.</p>
<p><b>Consult</b></p>	<p>No Prep, unless instructed otherwise</p>
<p><b>Coronary Artery Angiogram</b></p>	<p><b>Nothing to eat or drink for 6 hours prior to exam</b> (can have daily medications with a sip of water). No caffeine 12 hrs prior to exam. Must have a <b>driver</b>.</p>
<p><b>Coronary Artery Angiogram for A-fib</b></p>	<p>No Prep, unless instructed otherwise</p>
<p><b>CT Guided Cyst Aspiration</b></p>	<p>Please call 982-3477 to schedule a pre-admission appointment.  <b>Nothing to eat or drink for 6 hours prior to exam</b> (can have daily medications with a sip of water. <b>Withhold Coumadin for 5 days, &amp; Lovenox for 1 dose prior to exam after obtaining approval from prescribing physician.</b> Lab work as indicated at time of scheduling.</p>

	Must have a <b>driver</b> .
<b>CVC (Central Venous Catheter) Placement/Implant</b>	Please call 982-3477 to schedule a pre-admission appointment. <b>Nothing to eat or drink for 6 hours prior to exam</b> (can have daily medications with a sip of water). <b>Withhold Coumadin for 5 days, Lovenox for 1 dose, Plavix for 5 days, Pletal for 4 days, Pradaxa for 48 hours, &amp; Ticlid for 14 days prior to exam after obtaining approval from prescribing physician.</b> Lab work as indicated at time of scheduling. Must have a <b>driver</b> .
<b>Embolization, Neuro</b>	Please call 982-3477 to schedule a pre-admission appointment. <b>Nothing to eat or drink for 6 hours prior to exam</b> (can have daily medications with a sip of water). ( <b>Anesthesia patients</b> must have nothing to eat or drink for 8 hours prior to exam.) Lab work and bloodthinner restrictions as indicated at time of scheduling. Must have a <b>driver</b> .
<b>Embolization, Non-Neuro</b>	Please call 982-3477 to schedule a pre-admission appointment. <b>Nothing to eat or drink for 6 hours prior to exam</b> (can have daily medications with a sip of water). <b>Nothing to eat or drink for 6 hours prior to exam</b> (can have daily medications with a sip of water). <b>Withhold Coumadin for 5 days, Lovenox for 1 dose, Plavix for 5 days, Pletal for 4 days, Pradaxa for 48 hours, &amp; Ticlid for 14 days prior to exam after obtaining approval from prescribing physician.</b> Lab work as indicated at time of scheduling. ( <b>Anesthesia patients</b> must have nothing to eat or drink for 8 hours prior to exam.) Must have a <b>driver</b> .
<b>Epidural Steroid Injection</b>	No Prep, unless instructed otherwise
<b>Extremity Angiogram</b>	Please call 982-3477 to schedule a pre-admission appointment. <b>Nothing to eat or drink for 6 hours prior to exam</b> (can have daily medications with a sip of water). <b>Withhold Coumadin for 5 days, Lovenox for 1 dose, Plavix for 5 days, Pletal for 4 days, Pradaxa for 48 hours, &amp; Ticlid for 14 days prior to exam after obtaining approval from prescribing physician.</b> Lab work as indicated at time of scheduling. Must have a <b>driver</b> .
<b>Extremity Venogram</b>	Please call 982-3477 to schedule a pre-admission appointment. <b>Nothing to eat or drink for 6 hours prior to exam</b> (can have daily medications with a sip of water). <b>Withhold Coumadin for 5 days, &amp; Lovenox for 1 dose prior to exam after obtaining approval from prescribing physician.</b> Lab work as indicated at time of scheduling. Must have a <b>driver</b> .
<b>G or J Tube Exchange</b>	No Prep, unless instructed otherwise
<b>Gastrostomy Placement</b>	Please call 982-3477 to schedule a pre-admission

	<p>appointment.</p> <p><b>Nothing to eat or drink for 6 hours prior to exam</b> (can have daily medications with a sip of water). <b>Withhold Coumadin for 5 days, Lovenox for 1 dose, Plavix for 5 days, Pletal for 4 days, Pradaxa for 48 hours, &amp; Ticlid for 14 days prior to exam after obtaining approval from prescribing physician.</b> Lab work as indicated at time of scheduling. Must have a <b>driver</b>.</p>
<b>Hepatic Venogram with Hemodyn</b>	<p>Please call 982-3477 to schedule a pre-admission appointment.</p> <p><b>Nothing to eat or drink for 6 hours prior to exam</b> (can have daily medications with a sip of water). <b>Withhold Coumadin for 5 days &amp; Lovenox for 1 dose prior to exam after obtaining approval from prescribing physician.</b> Lab work as indicated at time of scheduling. Must have a <b>driver</b>.</p>
<b>IVC Filter Placement</b>	<p>Please call 982-3477 to schedule a pre-admission appointment.</p> <p><b>Nothing to eat or drink for 6 hours prior to exam</b> (can have daily medications with a sip of water). <b>Withhold Coumadin for 5 days &amp; Lovenox for 1 dose prior to exam after obtaining approval from prescribing physician.</b> Lab work as indicated at time of scheduling. Must have a <b>driver</b>.</p>
<b>Kyphoplasty Lumbar, Thoracic</b>	<p>Please call 982-3477 to schedule a pre-admission appointment.</p> <p><b>Nothing to eat or drink for 6 hours prior to exam</b> (can have daily medications with a sip of water). <b>Withhold Coumadin for 5 days, Lovenox for 1 dose, Plavix for 5 days, Pletal for 4 days, Pradaxa for 48 hours, &amp; Ticlid for 14 days prior to exam after obtaining approval from prescribing physician.</b> Lab work as indicated at time of scheduling. Must have a <b>driver</b>.</p>
<b>Liver Biopsy, Transjugular</b>	<p>Please call 982-3477 to schedule a pre-admission appointment.</p> <p><b>Nothing to eat or drink for 6 hours prior to exam</b> (can have daily medications with a sip of water). <b>Withhold Coumadin for 5 days &amp; Lovenox for 1 dose prior to exam after obtaining approval from prescribing physician.</b> Lab work as indicated at time of scheduling. Must have a <b>driver</b>.</p>
<b>Myelogram Cervical, Thoracic, Lumbar, Multilevel</b>	<p>Please call 982-3477 to schedule a pre-admission appointment.</p> <p><b>Nothing to eat or drink for 6 hours prior to exam</b> (can have daily medications with a sip of water). <b>Withhold Coumadin for 5 days, Lovenox for 1 dose, Plavix for 5 days, Pletal for 4 days, Pradaxa for 48 hours, &amp; Ticlid for 14 days prior to exam after obtaining approval from prescribing physician.</b> Lab work as indicated at time of scheduling. Must have a <b>driver</b>.</p>
<b>Nephrostomy Tube Change</b>	<p>Please call 982-3477 to schedule a pre-admission</p>

	<p>appointment.  <b>Nothing to eat or drink for 6 hours prior to exam</b> (can have daily medications with a sip of water). <b>Withhold Coumadin for 5 days &amp; Lovenox for 1 dose prior to exam after obtaining approval from prescribing physician.</b> Lab work as indicated at time of scheduling. Must have a <b>driver</b>.</p>
<p><b>Nephrostomy Tube Placement</b></p>	<p>Please call 982-3477 to schedule a pre-admission appointment.  <b>Nothing to eat or drink for 6 hours prior to exam</b> (can have daily medications with a sip of water). <b>Withhold Coumadin for 5 days, Lovenox for 1 dose, Plavix for 5 days, Pletal for 4 days, Pradaxa for 48 hours, &amp; Ticlid for 14 days prior to exam after obtaining approval from prescribing physician.</b> Lab work as indicated at time of scheduling. Must have a <b>driver</b>.</p>
<p><b>Nephrostomy Tube Removal</b></p>	<p>No Prep, unless instructed otherwise</p>
<p><b>Percutaneous Transhepatic Cholangiogram</b></p>	<p>Please call 982-3477 to schedule a pre-admission appointment.  <b>Nothing to eat or drink for 6 hours prior to exam</b> (can have daily medications with a sip of water). <b>Withhold Coumadin for 5 days, Lovenox for 1 dose, Plavix for 5 days, Pletal for 4 days, Pradaxa for 48 hours, &amp; Ticlid for 14 days prior to exam after obtaining approval from prescribing physician.</b> Lab work as indicated at time of scheduling. Must have a <b>driver</b>.</p>
<p><b>Pulmonary Angiogram</b></p>	<p>Please call 982-3477 to schedule a pre-admission appointment.  <b>Nothing to eat or drink for 6 hours prior to exam</b> (can have daily medications with a sip of water). <b>Withhold Coumadin for 5 days, Lovenox for 1 dose, Plavix for 5 days, Pletal for 4 days, Pradaxa for 48 hours, &amp; Ticlid for 14 days prior to exam after obtaining approval from prescribing physician.</b> Lab work as indicated at time of scheduling. Must have a <b>driver</b>.</p>
<p><b>Radio Frequency Ablation (RFA), Alcohol Ablation</b></p>	<p>Please call 982-3477 to schedule a pre-admission appointment.  <b>Nothing to eat or drink for 6 hours prior to exam</b> (can have daily medications with a sip of water). <b>Withhold Coumadin for 5 days, Lovenox for 1 dose, Plavix for 5 days, Pletal for 4 days, Pradaxa for 48 hours, &amp; Ticlid for 14 days prior to exam after obtaining approval from prescribing physician.</b> Lab work as indicated at time of scheduling. (<b>Anesthesia patients</b> must have nothing to eat or drink for 8 hours prior to exam.) Must have a <b>driver</b>.</p>
<p><b>Renal Angiogram</b></p>	<p>Please call 982-3477 to schedule a pre-admission appointment.  <b>Nothing to eat or drink for 6 hours prior to exam</b> (can have daily medications with a sip of water). <b>Withhold Coumadin for 5 days, Lovenox for 1 dose, Plavix for</b></p>

	<p><b>5 days, Pletal for 4 days, Pradaxa for 48 hours, &amp; Ticlid for 14 days prior to exam after obtaining approval from prescribing physician.</b> Lab work as indicated at time of scheduling. Must have a <b>driver</b>.</p>
<b>Renal Venogram</b>	<p>Please call 982-3477 to schedule a pre-admission appointment.  <b>Nothing to eat or drink for 6 hours prior to exam</b> (can have daily medications with a sip of water). <b>Withhold Coumadin for 5 days &amp; Lovenox for 1 dose prior to exam after obtaining approval from prescribing physician.</b> Lab work as indicated at time of scheduling. Must have a <b>driver</b>.</p>
<b>Sacroiliac Joint Injection</b>	<p>Please call 982-3477 to schedule a pre-admission appointment.  <b>Nothing to eat or drink for 6 hours prior to exam</b> (can have daily medications with a sip of water). <b>Withhold Coumadin for 5 days &amp; Lovenox for 1 dose prior to exam after obtaining approval from prescribing physician.</b> Lab work as indicated at time of scheduling. Must have a <b>driver</b>.</p>
<b>Spinal Angiogram/Arteriogram</b>	<p>Please call 982-3477 to schedule a pre-admission appointment.  <b>Nothing to eat or drink for 6 hours prior to exam</b> (can have daily medications with a sip of water). <b>Withhold Coumadin for 5 days &amp; Lovenox for 1 dose prior to exam after obtaining approval from prescribing physician.</b> Lab work as indicated at time of scheduling. Must have a <b>driver</b>.</p>
<b>Spinal Blood Patch</b>	<p>No Prep, unless instructed otherwise</p>
<b>Suprapubic Catheter Exchange</b>	<p>Please call 982-3477 to schedule a pre-admission appointment.  <b>Nothing to eat or drink for 6 hours prior to exam</b> (can have daily medications with a sip of water). <b>Withhold Coumadin for 5 days &amp; Lovenox for 1 dose prior to exam after obtaining approval from prescribing physician.</b> Lab work as indicated at time of scheduling. Must have a <b>driver</b>.</p>
<b>Suprapubic Catheter Placement</b>	<p>Please call 982-3477 to schedule a pre-admission appointment.  <b>Nothing to eat or drink for 6 hours prior to exam</b> (can have daily medications with a sip of water). <b>Withhold Coumadin for 5 days, Lovenox for 1 dose, Plavix for 5 days, Pletal for 4 days, Pradaxa for 48 hours, &amp; Ticlid for 14 days prior to exam after obtaining approval from prescribing physician.</b> Lab work as indicated at time of scheduling. Must have a <b>driver</b>.</p>
<b>SVC-Gram</b>	<p>Please call 982-3477 to schedule a pre-admission appointment.  <b>Nothing to eat or drink for 6 hours prior to exam</b> (can have daily medications with a sip of water). <b>Withhold Coumadin for 5 days, Lovenox for 1 dose, Plavix for</b></p>

	<b>5 days, Pletal for 4 days, Pradaxa for 48 hours, &amp; Ticlid for 14 days prior to exam after obtaining approval from prescribing physician.</b> Lab work as indicated at time of scheduling. Must have a <b>driver</b> .
<b>Thrombin Injection</b>	No Prep, unless instructed otherwise
<b>TIPS, TIPS Revision</b>	Please call 982-3477 to schedule a pre-admission appointment. <b>Nothing to eat or drink for 6 hours prior to exam</b> (can have daily medications with a sip of water). <b>Withhold Coumadin for 5 days, Lovenox for 1 dose, Plavix for 5 days, Pletal for 4 days, Pradaxa for 48 hours, &amp; Ticlid for 14 days prior to exam after obtaining approval from prescribing physician.</b> Lab work as indicated at time of scheduling. Must have a <b>driver</b> .
<b>Ultrasound Guided Cyst Aspiration</b>	Please call 982-3477 to schedule a pre-admission appointment. <b>Nothing to eat or drink for 6 hours prior to exam</b> (can have daily medications with a sip of water). <b>Withhold Coumadin for 5 days &amp; Lovenox for 1 dose prior to exam after obtaining approval from prescribing physician.</b> Lab work as indicated at time of scheduling. Must have a <b>driver</b> .
<b>Ureteral Stent Placement</b>	Please call 982-3477 to schedule a pre-admission appointment. <b>Nothing to eat or drink for 6 hours prior to exam</b> (can have daily medications with a sip of water). <b>Withhold Coumadin for 5 days, Lovenox for 1 dose, Plavix for 5 days, Pletal for 4 days, Pradaxa for 48 hours, &amp; Ticlid for 14 days prior to exam after obtaining approval from prescribing physician.</b> Lab work as indicated at time of scheduling. Must have a <b>driver</b> .
<b>Uterine Fibroid Embolization</b>	Please call 982-3477 to schedule a pre-admission appointment. <b>Nothing to eat or drink for 6 hours prior to exam</b> (can have daily medications with a sip of water). <b>Withhold Coumadin for 5 days, Lovenox for 1 dose, Plavix for 5 days, Pletal for 4 days, Pradaxa for 48 hours, &amp; Ticlid for 14 days prior to exam after obtaining approval from prescribing physician.</b> Lab work as indicated at time of scheduling. Must have a <b>driver</b> .
<b>Venogram Injection</b>	No Prep, unless instructed otherwise
<b>Venous Access Placement</b>	No Prep, unless instructed otherwise
<b>Vertebral Angiogram/Arteriogram</b>	Please call 982-3477 to schedule a pre-admission appointment. <b>Nothing to eat or drink for 6 hours prior to exam</b> (can have daily medications with a sip of water). <b>Withhold Coumadin for 5 days, Lovenox for 1 dose, Plavix for 5 days, Pletal for 4 days, Pradaxa for 48 hours, &amp; Ticlid for 14 days prior to exam after obtaining approval from prescribing physician.</b> Lab work as indicated at time of scheduling. Must have a <b>driver</b> .

<b>Vertebroplasty Thoracic, Lumbar</b>	Please call 982-3477 to schedule a pre-admission appointment. <b>Nothing to eat or drink for 6 hours prior to exam</b> (can have daily medications with a sip of water). <b>Withhold Coumadin for 5 days, Lovenox for 1 dose, Plavix for 5 days, Pletal for 4 days, Pradaxa for 48 hours, &amp; Ticlid for 14 days prior to exam after obtaining approval from prescribing physician.</b> Lab work as indicated at time of scheduling. Must have a <b>driver</b> .
<b>Visceral Angiogram/Arteriogram</b>	Please call 982-3477 to schedule a pre-admission appointment. <b>Nothing to eat or drink for 6 hours prior to exam</b> (can have daily medications with a sip of water). <b>Withhold Coumadin for 5 days, Lovenox for 1 dose, Plavix for 5 days, Pletal for 4 days, Pradaxa for 48 hours, &amp; Ticlid for 14 days prior to exam after obtaining approval from prescribing physician.</b> Lab work as indicated at time of scheduling. Must have a <b>driver</b> .